

SCHUYLER COUNTY HEALTH PROFILE

2017

ABOUT THE REPORT

The purpose of this report is to provide a summary of health data specific to Schuyler County. Where possible, benchmarks have been given to compare county rates to Upstate New York.

TOPIC	PAGE
About Schuyler County Population size, demographics, life expectancy, socioeconomic status and more	2-3
Health Behaviors Smoking status, diet, physical activity and chronic disease diagnoses	4-5
Social Determinants of Health Neighborhood accessibility, reported self-health and food insecurity	6
Life Expectancy Life expectancy at birth by ZIP code	7
Disparity A comparison of the highest and lowest estimated life expectancy ZIP codes by a number of health outcomes	8
Leading Causes of Death The five leading causes of death and premature death in Schuyler County	9
Population Health Measures Sixteen health metrics selected to measure community impact on population health for Schuyler County	10-11
End Notes Information on data sources used throughout the report	12-14
About Common Ground Health Who we are	15

Where comparisons between regions are made throughout the report, the following color code will be used:




About Schuyler County

18,458 RESIDENTS LIVE IN SCHUYLER COUNTY.

17,660 are White Non-Hispanic.



POPULATION BY AGE GROUP



33.9% of adults reported experiencing housing insecurity in the past 12 months.

4.6% of the labor force is unemployed.


13.7% of the population is living with a disability.

77.9 years is the average life expectancy at birth.

About Schuyler County

More than 2,800 residents – 15.4 percent of Schuyler County’s population – live below the federal poverty level according to U.S. Census statistics. As the map illustrates, the concentration of poverty is highest in Dundee, Bradford and Montour Falls. Research shows that lower socioeconomic status (SES) is linked to higher incidence of chronic disease, shorter life expectancy and lower rates of good social, emotional and physical health.¹

SES BY ZIP CODE - Schuyler County



Education levels can also predict life expectancy. The Centers for Disease Control and Prevention reports that adults aged 25 without a high school diploma “can expect to die nine years sooner than college graduates.”² Approximately 50 percent of Schuyler County residents have at least some form of college experience, compared to 59 percent of New York state residents.

	High school graduate or less, 50%	College experience, 32%	Bachelors degree or higher, 18%
Schuyler County			
NYS	High school graduate or less, 42%	College experience, 25%	Bachelors degree or higher, 34%

Health Behaviors

Behavioral and personal lifestyles are important determinants of health. Smoking, poor nutrition and other unhealthy behaviors are linked to adverse health outcomes.

Many of the indicators for behavioral lifestyles shown below for Schuyler County are similar to Upstate New York Rates. However, there is a lower percent of persons participating in leisure time physical activity and consumption of sugary beverages in Schuyler County.




	SCHUYLER COUNTY	UPSTATE NEW YORK
Consumed fast food three or more times per week	5.5%	6.3%
Consumed one or more sugary drinks daily	18.9%	23.3%
Participated in leisure time physical activity in the past 30 days	67.6%	73.7%

Adult smoking rates are higher in Schuyler County (22.3 percent) compared to Upstate New York (17.3 percent). Data reveal an estimate of more than 3,200 adults who currently smoke in Schuyler County.

	SCHUYLER COUNTY	UPSTATE NEW YORK
Current cigarette smoker	22.3%	17.3%


Health Behaviors

Inactive lifestyles and poor diets may lead to obesity, a risk factor for developing diabetes, hypertension and other chronic illnesses. Estimated rates of obese or overweight adults, hypertension and diabetes in Schuyler County are similar to Upstate New York. However, there is a slightly higher prevalence of hypertension in Schuyler County (32.5 percent).

	SCHUYLER COUNTY	UPSTATE NEW YORK
 Obese or overweight adults	59.2%	62.2%
 Physician-diagnosed diabetes	10.6%	9.2%
 Physician-diagnosed hypertension	32.5%	30.2%

For individuals with hypertension, controlling high blood pressure with medication and lifestyle change is critical to avoiding complications such as heart attack, stroke and kidney failure. The individuals who are most at risk for these life threatening events are those who have extremely high blood pressure, known as stage 2 hypertension. As of June 2016, the region's high blood pressure registry showed that more than 71 percent of Schuyler County adults with hypertension had their condition under control. For residents whose blood pressure was uncontrolled, 15 percent had stage 2 hypertension.



Uncontrolled Hypertension by Status - June 2016



Social Determinants of Health

The physical environment plays an important role in residents' ability to engage in physical activity and access nutritious food. Many residents in Schuyler County experience indicators of poor environmental health, including experiencing food insecurity.

Of note, Schuyler County has a slightly higher percent of its population reporting food insecurity in the past year (24.0 percent) compared to Upstate New York (22.7 percent).

	SCHUYLER COUNTY	UPSTATE NEW YORK
 Consider neighborhood suitable for walking and physical activity	94.7%	92.3%
 Reported food insecurity in the past 12 months	24.0%	22.7%

Approximately one in six (16.9 percent) Schuyler County residents report experiencing 14 or more poor mental health days the past month. This is compared to 11.5 percent of Upstate New York residents.




In addition, over 1,800 Schuyler County residents reported 14 or more poor days of physical health (12.6 percent), similar to Upstate New York rates (12.3 percent).

Life Expectancy

Although average life expectancy in Schuyler County is 77.9 years, how long residents live on average varies by more than 3 years depending on their ZIP code.

For example, ZIP code area 14865, Montour Falls, has an estimated life expectancy of only 75.2 years at birth. By contrast, the county's highest estimated life expectancy, 78.5 years, is found in Beaver Dams 14812 ZIP code area.

LIFE EXPECTANCY BY ZIP CODE - Schuyler County




Research has shown that poverty is associated with shorter life expectancy. Residents with lower socioeconomic status are less likely to seek preventative care and to monitor/maintain good health behaviors for a variety of reasons.

Disparity

A comparison of ZIP codes shows that rates for negative health indicators, including rates of Prevention Quality Indicators* and premature death are higher in Montour Falls than in Beaver Dams. In addition, there are significantly higher rates of teenage pregnancy in Montour Falls.

As some of the selected indicators reveal unexpected data for these two ZIP codes, there are likely other factors not captured here that are driving the disparity in life expectancies.




	14865 - MONTOUR FALLS	14812 - BEAVER DAMS
Prevention Quality Indicators (Adult)*	3,318 per 100,000 adults	1,857 per 100,000 adults
Premature Death (before age 75)	4,656 per 100,000 population	3,397 per 100,000 population
Premature Births	1.3% of births	11.5% of births
Late or No Prenatal Care	2.3% of births	4.4% of births
Teenage Pregnancy Rate	63.7 per 1,000 population	35.4 per 1,000 population

*Prevention Quality Indicators (PQI) were developed by the Agency for Healthcare Research and Quality and are based on ICD-9 coding and classifies conditions that are seen as potentially preventable/avoidable had sufficient care been given in the primary care setting prior to the presentation.


Leading Causes of Death

Population Health Measures

Leading causes of death in Schuyler County are similar those of Upstate New York. However, rates of mortality from cancer are higher in Schuyler County than in Upstate New York. The graph below demonstrates the five leading causes of death.



Many of the leading causes of death shown in the figure above can be attributed to chronic diseases such as hypertension and obesity. However, the data below tell a slightly different story for premature death before age 75. Unintentional injury and suicide rank among the top five causes of premature death in Schuyler County.



In 2013, Common Ground Health convened the Regional Commission on Community Health Improvement. Over 18 months, the Commission engaged leaders from across the region to study and develop ideas for improving the integration of care and addressing the complex medical, behavioral and social needs of vulnerable community residents.

The Commission adopted a set of communitywide measures to gauge the region's collective progress toward improved community health. Below are the population health measures selected by the Commission specific to Schuyler County.

REGIONAL TREND INDICATOR:

The icons indicate whether the Finger Lakes region is getting better, has remained flat or is getting worse for each measure.












COMMUNITY MEASURES	SCHUYLER COUNTY	FINGER LAKES REGION
Childhood Immunization Percentage of children receiving recommended immunizations by 36 months	64.3%	64.5%
Childhood Obesity Percentage of school children whose weight is at the 95th percentile or greater	16.1%	16.1%
Adult Obesity Percentage of adults 18 years or older with a BMI of 30 or greater	24.1%	27.0%
Adult Smoking Percentage of adults who are current smokers	22.3%	17.4%
High Blood Pressure Control Percentage of adults with high blood pressure who had their condition under control	71.0%	68.0%
Type II Diabetes Control Under development	Under Development	Under Development
High School Graduation Percentage of students graduating in four years	88.0%	81.2%

Population Health Measures

End Notes



 CLINICAL MEASURES	SCHUYLER COUNTY	FINGER LAKES REGION
Preventable Hospital Stays (rate per 100,000 population) Number of hospital stays for PQIs	1,443	1,074 
Depression Screening Under development	Under Development	Under Development
ED Visits with Behavioral Health Diagnosis Percentage of all ED patients with a behavioral health diagnosis	5.4%	8.6% 
ED Visits Among those 65+ (rate per 1,000 population) Number of ED visits that involve adults 65+	539	303 
30 Day Readmissions with Behavioral Health Diagnoses 30-day behavioral health-related hospital readmission rates	12.3%	13.4% 
Nursing Home Use Number of nursing home days per 1,000 individuals 85+	108,163	96,053

 HEALTH OUTCOMES	SCHUYLER COUNTY	FINGER LAKES REGION
Premature Death (rate per 100,000 population) Years of potential life lost before age 65	2,416	3,496 
Low Birth Weight Percentage of births with birthweight less than 2500 g/ 5.5 lbs	9.2%	7.8% 
Good Health Self-Report Percentage of adults reporting excellent, very good or good general health status	85.0%	83.7% 

References

1. **The Future of Children:** A collaboration of the Woodrow Wilson School of Public and International Affairs at Princeton University and The Brookings Institution. "The Health-Related Effects of Socioeconomic Status," February 2013.
2. **Health, United States, 2011:** with special feature on socioeconomic status and health. Hyattsville, MD: National Center for Health Statistics; 2012.

Sources for Population Health Measures

Community Measures

- Childhood Immunization** – New York State Immunization Information System, 2013
- Childhood Obesity** – Student Weight Status Category Reporting System, 2012-2014
- Adult Obesity** – Expanded Behavioral Risk Factor Surveillance System (ExpBRFSS) 2013-2014
- Adult Smoking** – ExpBRFSS 2013-2014
- High Blood Pressure Control** – Common Ground Health Hypertension Registry, June 2016
- Type II Diabetes Control** – Under Development
- High School Graduation** – NYS Department of Education, 2015

Clinical Measures

- Preventable Hospital Stays** – NY Statewide Planning and Research Cooperative System (SPARCS), 2014-2015
- Depression Screening** – Under Development
- ED Visits with a Behavioral Health Diagnosis** – SPARCS, 2014
- ED Visits among those 65+** – SPARCS, 2014
- 30 Day Readmissions with Behavioral Health Diagnoses** – SPARCS, 2014
- Nursing Home Use among 85+** – Centers for Medicare and Medicaid Skilled Nursing Facility Cost Reports, 2013

Health Outcomes

- Premature Death** – NYS Vital Statistics, 2014
- Low Birthweight** – NYS Perinatal Data Profile, 2015
- Good Health Self-Report** – ExpBRFSS, 2013-2014

Data Sources

U.S. Census Bureau/American Community Survey: Although the American Community Survey (ACS) produces population, demographic and housing unit estimates, it is the Census Bureau's Population Estimates Program that produces and disseminates the official estimates of the population for the nation, states, counties, cities and towns and estimates of housing units for states and counties. Data are based on a sample and are subject to sampling variability. The value shown here is the 90 percent margin of error. Estimates of urban and rural population, housing units, and characteristics reflect boundaries of urban areas defined based on Census 2010 data. As a result, data for urban and rural areas from the ACS do not necessarily reflect the results of ongoing urbanization.

End Notes

New York State Department of Labor: Civilian Unemployment includes those individuals who were not working but were able, available and actively looking for work during the week including the 12th of the month. Individuals who were waiting to be recalled from a layoff, and individuals waiting to report to a new job within 30 days were also considered to be unemployed. Unemployment Rate is the number of unemployed as a percentage of the labor force.

Expanded Behavioral Risk Factor Surveillance System/Sub-County Health Data Report: Data was collected for ExpBRFSS over the course of 12 monthly waves, starting April 15, 2013 and ending May 10, 2014. The goal of each wave was to obtain roughly 8.3 percent of the required completes in each county for the landline survey and each region for the cell phone survey. Experienced interviewers conducted telephone interviews using computer-assisted telephone interviewing (CATI) software.

Common Ground Health High Blood Pressure Registry: The Common Ground Health hypertension registry collects data from medical practices and systems on hypertensive patients. Data provided are reflective of the patients involved in the registry. The Healthcare Effectiveness Data and Information Set (HEDIS) is the tool used to measure hypertension control. Controlled hypertension are those who have a reading in the past year <140/90; Stage 1 are those with a reading between 141/90—159/99; Stage 2 are those with a reading >160/100. Those who have not had a reading in the past 12 months are considered uncontrolled.

United States Department of Agriculture Economic Research Service: Low access to healthy food is defined as being far from a supermarket, supercenter or large grocery store ("supermarket" for short). Food desert status is defined as a low income census tract with a supermarket more than a half of a mile away for urban tracts, and more than 10 miles away for rural tracts. Population size is according to the 2010 Census estimates. A census tract is considered to have low access if a significant number or share of individuals in the tract is far from a supermarket. A significant number of the population includes at least 500 people or 33 percent of the tract's population.

New York State Department of Health Vital Statistics: The cause of death reported in this publication is the underlying cause classified according to the tenth revision of the International Classification of Diseases (ICD, 10th revision) adopted by New York state in 1999. Historically, several revisions of the ICD have been used, therefore, it is necessary to employ a comparability ratio when comparing cause of death statistics across revisions. Comparability ratios have been published by the National Center for Health Statistics (NCHS).

New York State Perinatal Data Profile: Premature births are those which occurred prior to 37 weeks gestation. Low birthweight babies are births weighing between 100-2499 grams. The teenage pregnancy rate looks at pregnancies (births + abortions + spontaneous fetal deaths) to females ages 15-19 per 1,000 female population ages 15-19. Rates are computed using the three-year average number of teen pregnancies and the population for the middle year of the three-year time period.

Statewide Planning and Research Cooperative System: SPARCS is a comprehensive all payer data reporting system established in 1979 as a result of cooperation between the healthcare industry and government. The system was initially created to collect information on discharges from hospitals. SPARCS currently collects patient level detail on patient characteristics, diagnoses and treatments, services, and charges for each hospital inpatient stay and outpatient (ambulatory surgery, emergency department and outpatient services) visit; and each ambulatory surgery and outpatient services visit to a hospital extension clinic and diagnostic and treatment center licensed to provide ambulatory surgery services. All calculations were performed by Common Ground Health and are age/sex adjusted rates.

New York State Immunization Information System (NYSIIS): NYSIIS is a statewide immunization information system or registry which maintains immunization data of persons of all ages. The recommended vaccinations by 36 months of age include the 4:3:1:3:3:1:4 immunization series: 4DTaP, 3 polio, 1 MMR, 3 hep B, 3 Hib, 1 varicella, and 4 PCV13.


Student Weight Status Category Reporting System (SWSCR): The SWSCR collects data on weight status category, including underweight, healthy weight, overweight or obese based on BMI-for-age percentile on children and adolescents attending public schools in New York state.


NYS Department of Education: Data are submitted to the New York State Department of Education (NYSED) by school districts, charter schools, Boards of Cooperative Educational Services (BOCES), institutions of Higher Education and NYSED program offices.

Centers for Medicare and Medicaid Skilled Nursing Facility Cost Reports: Data are collected on nursing home days from 2013 residential health care facility cost reports. Population estimates are collected from the U.S. Census Bureau, June 2015.

For reports on the remaining 8 counties in the Finger Lakes Region, and additional data reports for Schuyler County, visit: www.commongroundhealth.org/countyprofiles

For questions regarding this report, contact Catie Kunecki at: Catie.Kunecki@commongroundhealth.org or (585) 224-3157





ABOUT COMMON GROUND HEALTH

Founded in 1974, Common Ground Health is one of the nation's oldest and most effective regional health planning organizations. Located in Rochester, N.Y., the nonprofit serves the nine-county Finger Lakes region. We bring together health care, education, business, government and other sectors to find common ground on health issues. Learn more about our community tables, our data resources and our work improving population health at www.CommonGroundHealth.org.

1150 University Avenue | Rochester NY 14607
585.224.3101 | CommonGroundHealth.org

